New Statistical Techniques for Predictive Water Quality Modeling at Great Lakes Beaches



Michael N. Fienen¹
Wesley R. Brooks¹
Steven R. Corsi¹
Kurt Wolfe²
Rajbir Parmar²
Mike Galvin²
Mike Cyterski²



¹USGS Wisconsin Water Science Center, Middleton, WI ²USEPA Office of Research and Development, Athens, GA

Introduction

E. coli is used as a fecal indicator bacterium

Without rapid methods, 24 hours are required to get an answer

Environmental predictors (weather, waves, etc.) are available in real time

Our ultimate goal is to predict *E. coli* from environmental predictors.





Modeling Approaches

getting the most information from all data available

persistence model

using yesterday's *E. coli* to predict today's conditions

ordinary least squares regression (OLS)

well-established technique

requires decisions on how to handle interaction and correlation

partial least squares (PLS)

newer technique used extensively in spectroscopy

introduced to beach community by Hou, Rabinovici, and Boehm (2006)


useful when variables are correlated or insensitive

overfitting prevented through cross validation and component trimming

algorithm replaces trial-and-error interaction terms & variable selection



Partial Least Squares: Example





Partial Least Squares

regression built on principal directions relating variables to predictions


$$\mathbf{y} = \mathbf{X}\boldsymbol{\beta} + \boldsymbol{\mathcal{E}}$$
 error term coefficients observations predictor variables

principal directions of covariance between **X** and **y** define components

using all components is equivalent to OLS with all variables


the number of components retained is chosen balancing lower PRESS against overfitting

each component includes information from all base variables





Decision and Regulatory Thresholds






Tradeoffs for Model Performance

robust methods and improved data integration

building a PLS model

managers decide the tradeoff between protective and permissive



sensitivity: proportion of true positives

high sensitivity means increasing true positives

specificity: proportion of true negatives

low specificity means increasing false positives


specificity limit (vertical blue line) is the dial controlling this tradeoff




Great Lakes Beaches Modeled

Wisconsin Department of Natural Resources









Building a PLS model – example for Edgewater, Ohio

2005-2010: split training data


Build and test models: predict each fold using data from the other four

2011: prediction year

divide the training data randomly into five equallysized "folds" select a few candidate specificity limits (tuning) compare model performance on the test folds pick the specificity limit that had the best performance, and train a new model over all five folds make and record
predictions to manage
beach and provide
data going forward




2010 Model Performance






Next Steps



Next Steps

Virtual Beach

Well-known EPA software for predicting bacterial concentration PLS regression to be included in version 3 (coming 2012)





Acknowledgements

OLS model results and collaboration:

Rob Darner

Donna Francy

Amie Brady

Dan Ziegler

Adam Mednick

Data provided by:

Cuyahoga County Board of Health

Northeast Ohio Regional Sewer District

Ozaukee County, Wisconsin

Funding provided by:

Ocean Research Priorities Plan (ORPP)

USEPA Great Lakes Restoration Initiative (GLRI)



